

Registration for the Spring-Meeting of the  
German Physical Society  
from 04.03. to 09.03.2005  
in berlinamop

**Passively mode-locked laser performance of Yb:KLuW** —  
•SIMON RIVIER<sup>1</sup>, XAVIER MATEOS<sup>1</sup>, VALENTIN PETROV<sup>1</sup>, UWE  
GRIEBNER<sup>1</sup>, MARTIN ZORN<sup>2</sup>, and MARKUS WEYERS<sup>2</sup> — <sup>1</sup>Max-Born-  
Institut, Max-Born-Str. 2a, 12489 Berlin — <sup>2</sup>Ferdinand-Braun-Institut,  
Albert-Einstein-Str. 11, 12489 Berlin

Yb-doped laser materials have received much attention for the generation of ultrashort pulses. One of the most promising results have been obtained with the monoclinic double tungstates Yb:KGW and Yb:KYW [1]. The novel Yb-doped tungstate host KLuW [2] was investigated applying diode- and Ti:Sa-pumped laser schemes. The mode-locked laser performance of Yb:KLuW was compared in the pico- and femtosecond regime for the polarization parallel to the crystallo-optic Nm- and Np-axis. Using end-pumped configurations and a semiconductor saturable absorber (SAM) in a standard type-z laser cavity, laser operation in the 100fs-range with an average power up to 300mW around 1045nm was achieved. Pulses as short as 81fs duration could be generated at a repetition rate of 94MHz with a time-bandwidth product of 0.32. The demonstrated pulse durations are the shortest obtained with Yb-doped tungstate-based oscillators mode-locked with SAM.

[1] P. Klopp et al., Opt. Express 10, 108 (2002). [2] X. Mateos et al., IEEE J. Quantum. Electron. 40, 1056 (2004).

Location: berlinamop  
Date: 04.03.—09.03.2005  
Section: Short Time-scale Physics  
Subject: laser systems  
Presentation: Talk  
Email: rivier@mbi-berlin.de  
Membership: none